

Spring 16 – AMS256 Homework 1

1. Observe that $z_i = y_i/x_i = \beta_1 + \beta_0/x_i + \epsilon_i/x_i \Rightarrow$ This model is still linear in β .
2. (a) Note that three columns are linearly independent. Or the three rows are linearly independent. So $r(\mathbf{X}) = 3$
- (b) $\dim(C(\mathbf{X})) = r(\mathbf{X}) = 3$.

A basis that we can easily come up with (you may find a different set of vectors for the

basis) is a set of the following three vectors, $\mathbf{u}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$ or $\mathbf{u}_2 = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}$ or $\mathbf{u}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}$.

- (c) $\dim(\mathcal{N}(\mathbf{X})) = 4 - \dim(C(\mathbf{X})) = 4 - 3 = 1$. We need to find a vector \mathbf{v} such that

$\mathbf{X}\mathbf{v} = \mathbf{0}$. A basis is $\mathbf{v}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}$. We can check that $\mathbf{v}_1 \perp \mathbf{u}_j, j = 1, 2, 3$.

- (d) $\mathbf{X}^T\mathbf{X} = \begin{bmatrix} 5 & 1 & 3 & 1 \\ 1 & 3 & 1 & 3 \\ 3 & 1 & 3 & 1 \\ 1 & 3 & 1 & 3 \end{bmatrix}$. Recognize that the submatrix of the upper 3×3 is nonsingular

matrix and find a g-inverse matrix, $(\mathbf{X}^T\mathbf{X})^- = \begin{bmatrix} 1/2 & 0 & -1/2 & 0 \\ 0 & 3/8 & -1/8 & 0 \\ -1/2 & -1/8 & 7/8 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

- (e)

$$\mathbf{P} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^-\mathbf{X}^T = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1/2 & 1/2 & 0 & 0 \\ 0 & 1/2 & 1/2 & 0 & 0 \\ 0 & 0 & 0 & 1/2 & 1/2 \\ 0 & 0 & 0 & 1/2 & 1/2 \end{bmatrix}$$

If $\mathbf{y} \in C(\mathbf{P}) = C(\mathbf{X})$, $\mathbf{P}\mathbf{y} = \mathbf{y}$. So $[3, 1, 1, 2, 2]^T$ and $[1, 0, 0, 2, 2]^T$ can be $\mathbf{P}\mathbf{y}$.

- (f) Recall that $(\mathbf{X}^T\mathbf{X})^-\mathbf{X}$ is a g-inverse of \mathbf{X} . We use this to construct all the solutions, $\tilde{\mathbf{x}} = \mathbf{G}\mathbf{c} + (\mathbf{I} - \mathbf{G}\mathbf{A})\mathbf{z}$ for $\mathbf{A}\mathbf{x} = \mathbf{c}$ from lecture. Thus, for some $\mathbf{z} \in \mathbb{R}^p$

$$\tilde{\mathbf{x}} = (\mathbf{X}^T\mathbf{X})^-(\mathbf{X}^T\mathbf{X})\mathbf{y} + (\mathbf{I} - (\mathbf{X}^T\mathbf{X})^-(\mathbf{X}^T\mathbf{X}))\mathbf{z}$$

By varying \mathbf{z} , we can obtain all possible solutions of β .

- 3.

$$\begin{aligned} y_i &= \beta_0 + \beta_1 x_i + \epsilon_i \\ &= \beta_0 + \beta_1(s + ti) + \epsilon_i \\ &= \beta_0 + \beta_1 s + \beta_1 ti + \epsilon_i \\ &= \gamma_0 + \gamma_1 i + \epsilon_i \end{aligned}$$

where $\gamma_0 = \beta_0 + \beta_1 s$ and $\gamma_1 = \beta_1 t$. Thus, they are an equivalent parameterization.

4. • LSE

Let $\hat{\theta} = \operatorname{argmin}_{\theta} Q(\theta) = \operatorname{argmin}_{\theta} \|\mathbf{Y} - \mathbf{X}\beta\|$ where $\mathbf{X} = [x_1^2, \dots, x_n^2]^T$ and $\beta = \theta$. From lecture, for full rank \mathbf{X} we know

$$\hat{\beta} = \hat{\theta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y} = \frac{\sum_i x_i^2 y_i}{x_i^4}.$$

• MLE

$$\frac{\partial \log f(\mathbf{y}|\theta)}{\partial \theta} = \frac{\partial \left\{ -\frac{n}{2} \log(2\pi\sigma^2) - \frac{\sum_i (y_i - \theta x_i)^2}{2\sigma^2} \right\}}{\partial \theta} = 0 \Rightarrow \hat{\theta} = \frac{\sum_{i=1}^n x_i^2 y_i}{\sum_{i=1}^n x_i^4}$$

5. Plugging in \mathbf{x}^* and \mathbf{y}^* for \mathbf{x} and \mathbf{y} , respectively, we obtain

$$\begin{aligned} \hat{\beta}_1^* &= \frac{S_{xy}^*}{S_{xx}^*} = \frac{\sum (x_i^* - \bar{x}^*)(y_i^* - \bar{y}^*)}{\sum (x_i^* - \bar{x}^*)^2} = \frac{\sum ((c + dx_i) - (c + d\bar{x}))((a + by_i) - (a + b\bar{y}))}{\sum ((c + dx_i) - (c + d\bar{x}))^2} \\ &= \frac{b \sum (x_i - \bar{x})(y_i - \bar{y})}{d \sum (x_i - \bar{x})^2} = \frac{b\hat{\beta}_1}{d}, \\ \hat{\beta}_0^* &= \bar{y}^* - \hat{\beta}_1^* \bar{x}^* = (a + b\bar{y}) - \frac{b\hat{\beta}_1}{d}(c + d\bar{x}) = a - \frac{bc\hat{\beta}_1}{d} + b\hat{\beta}_0. \end{aligned}$$

6. From lecture, $\hat{\beta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$.

$$\mathbf{X}^T \mathbf{X} = \begin{bmatrix} 8 & 272 \\ 272 & 9318 \end{bmatrix} \Rightarrow (\mathbf{X}^T \mathbf{X})^{-1} = \begin{bmatrix} 16.639 & -0.486 \\ -0.486 & 0.014 \end{bmatrix}$$

$$\mathbf{X}^T \mathbf{Y} = \begin{bmatrix} 71 & 63 & 68 & 70 & 71 & 63 & 68 & 70 \\ 2457 & 2010 & 2176 & 2176 & 2205 & 2412 & 2448 & 1920 \end{bmatrix}^T$$

$$\Rightarrow \hat{\beta} = \begin{bmatrix} 35.174 \\ 0.929 \end{bmatrix}$$

ANOVA Table:

Source of Variation	Sum of Squares	DF	Mean squares	F-stat
Model	60.415	1	60.415	50.73
Error(residual)	7.145	6	1.19	
Total	67.5	7	9.64	

$$\begin{aligned} \sum_{i=1}^n (\hat{y}_i - \bar{y})^2 &= \sum_{i=1}^n (\hat{\beta}_0 + \hat{\beta}_1 x_i - \hat{\beta}_0 - \hat{\beta}_1 \bar{x})^2 = \hat{\beta}_1^2 \sum_{i=1}^n (x_i - \bar{x})^2 \\ &= \left(\frac{S_{xy}}{S_{xx}} \right)^2 \sum_{i=1}^n (x_i - \bar{x})^2 = \frac{S_{xy}^2}{S_{xx}} = \mathbf{y}^T (\mathbf{P}_x - \mathbf{P}_1) \mathbf{y}, \end{aligned}$$

where \mathbf{P}_x and \mathbf{P}_1 are the orthogonal projection operators onto $C(\mathbf{X})$ and $C(\mathbf{1})$, respectively. That is, \mathbf{P}_x : projection matrix for the model for an intercept and a slope and \mathbf{P}_1 : projection matrix under the model with an intercept. From lecture,


$$R^2 = 1 - \frac{\|\mathbf{y}^T(\mathbf{P}_x - \mathbf{P}_1)\mathbf{y}\|^2}{\|\mathbf{y}^T(\mathbf{I} - \mathbf{P}_1)\mathbf{y}\|^2} = 1 - \frac{\sum_i(\hat{y}_i - \bar{y})^2}{\sum_i(y_i - \bar{y})^2} = 1 - \frac{SSE}{SST} = 1 - \frac{7.145}{67.5} = 0.894$$

7. (a)

```
modell=lm(MPG~HP, dat)
plot(dat$HP, dat$MPG)
abline(modell)
summary(modell)
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	50.0661	1.5695	31.90	0.0000
HP	-0.1390	0.0121	-11.52	0.0000


Residual standard error: 6.174 on 80 degrees of freedom
 Multiple R-squared: 0.6239,
 Adjusted R-squared: 0.6192
 F-statistic: 132.7 on 1 and 80 DF,
 p-value: < 2.2e-16



```
(b) model2=lm(log(MPG)~HP, dat)
plot(dat$HP, log(dat$MPG))
abline(model2)
summary(model2)
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.0132	0.0401	100.02	0.0000
HP	-0.0046	0.0003	-14.87	0.0000

Residual standard error: 0.1578 on 80 degrees of freedom
Multiple R-squared: 0.7344,
Adjusted R-squared: 0.7311
F-statistic: 221.2 on 1 and 80 DF, p-value: < 2.2e-16



The scatterplots shows that the linear model fits better for $\log(MPG)$. Also, from the R output, Model2 has a higher Adjusted R-squared compared to Model1, which indicates that Model2 has better explains the variability in the data better than Model1. Model2 also have a lower residual standard error compared to Model1.

```
(c) model3=lm(MPG~HP+VOL+SP+WT, dat)
summary(model3)
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	192.4378	23.5316	8.18	0.0000
HP	0.3922	0.0814	4.82	0.0000
VOL	-0.0156	0.0228	-0.69	0.4951
SP	-1.2948	0.2448	-5.29	0.0000
WT	-1.8598	0.2134	-8.72	0.0000

Residual standard error: 3.653 on 77 degrees of freedom
Multiple R-squared: 0.8733,
Adjusted R-squared: 0.8667
F-statistic: 132.7 on 4 and 77 DF, p-value: < 2.2e-16

8.

$$\begin{aligned} E(y_0) &= E(\hat{\beta}_0 + \hat{\beta}_1 x_0 + \epsilon_0) \\ &= \beta_0 + \beta_1 x_0 \end{aligned}$$

$$\begin{aligned} \text{Var}(y_0) &= \text{Var}(\hat{\beta}_0 + \hat{\beta}_1 x_0 + \epsilon_0) \\ &= \text{Var}(\hat{\beta}_0 + \hat{\beta}_1 x_0) + \text{Var}(\epsilon_0) \\ &= \text{Var}(\hat{\beta}_0 + \hat{\beta}_1 x_0) + \sigma^2 \\ &= \begin{bmatrix} 1 & x_0 \end{bmatrix} \begin{bmatrix} \frac{\sigma^2}{nS_{xx}} \sum x_i^2 & -\frac{\sigma^2 \bar{x}}{nS_{xx}} \\ -\frac{\sigma^2 \bar{x}}{nS_{xx}} & \frac{\sigma^2}{nS_{xx}} \end{bmatrix} \begin{bmatrix} 1 \\ x_0 \end{bmatrix} + \sigma^2 \\ &= \frac{\sigma^2}{nS_{xx}} \left(\frac{\sum x_i^2}{n} - 2\bar{x}x_0 + x_0^2 \right) + \sigma^2 \end{aligned}$$